



TITLE:

NORMAL INTERMEDIATE SUBFACTORS

AUTHOR(S):

TERUYA, TAMOTSU

CITATION:

TERUYA, TAMOTSU. NORMAL INTERMEDIATE SUBFACTORS. 数理解析研
究所講究録 1996, 956: 90-115

ISSUE DATE:

1996-08

URL:

<http://hdl.handle.net/2433/60425>

RIGHT:

NORMAL INTERMEDIATE SUBFACTORS

TAMOTSU TERUYA

DEPARTMENT OF MATHEMATICS

HOKKAIDO UNIVERSITY

SAPPORO 060 JAPAN

1. INTRODUCTION

It is well known that if $\alpha : G \rightarrow \text{Aut}(N)$ is an outer action of a finite group G on a type II_1 factor N and K is an intermediate subfactor for $N \subset N \rtimes_{\alpha} G$, then there is a subgroup H of G such that $K = N \rtimes_{\alpha} H$ (see for instance [19]). On the other hand, Y. Watatani [31] showed that there exist only finitely many intermediate subfactors for an irreducible inclusion with finite index. So it is natural to consider intermediate subfactors as “quantized subgroups” in the index theory for an inclusion of type II_1 factors. The notion of normality for subgroups plays important role in the theory of finite groups. In this paper we introduce the notion of normality for intermediate subfactors of irreducible inclusions.

D. Bisch [1] and A. Ocneanu [21] gave a nice characterization of intermediate subfactors of a given irreducible inclusion $N \subset M$ in terms of Jones projections and Ocneanu’s Fourier transform $\mathcal{F} : N' \cap M_1 \rightarrow M' \cap M_2$. We define normal intermediate

subfactors as follows:

Definition. Let $N \subset M$ be an irreducible inclusion of type II_1 factors with finite index and K an intermediate subfactor of the inclusion $N \subset M$. Then K is a *normal intermediate subfactor* of the inclusion $N \subset M$ if $e_K \in \mathcal{Z}(N' \cap M_1)$ and $\mathcal{F}(e_K) \in \mathcal{Z}(M' \cap M_2)$, where e_K is the Jones projection for the inclusion $K \subset M$.

Every finite dimensional Hopf C^* -algebra (Kac algebra) gives rise to an irreducible inclusion of AFD II_1 factors, which are characterized by depth 2 (see for example [21], [28], [29], [34]). Let M be the crossed product algebra $N \rtimes \mathbf{H}$ of N by an outer action of a finite dimensional Hopf C^* -algebra \mathbf{H} . Unfortunately, there is no one-to-one correspondence between the intermediate subfactors of $N \subset M$ and the subHopf C^* -algebras of \mathbf{H} in general. But we get the next result:

Theorem. Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index, i.e., M is described as the crossed product algebra $N \rtimes \mathbf{H}$ of N by an outer action of a finite dimensional Hopf C^* -algebra \mathbf{H} . Let K be an intermediate subfactor of $N \subset M$ and e_K is the Jones projection for $K \subset M$. Then K is described as the crossed product algebra $N \rtimes \mathbf{K}$ of N by an outer action of a subHopf C^* algebra \mathbf{K} of \mathbf{H} if and only if e_K is an element of the center of the relative commutant algebra $N' \cap M_1$, where M_1 is the basic extension for $N \subset M$.

Let $N \subset M$ be an irreducible inclusion of type II_1 factors with finite index and M_1 the basic extension for $N \subset M$. Let K be an intermediate subfactor of $N \subset M$

and K_1 the basic extension for $K \subset M$. Then K_1 is an intermediate subfactor of $M \subset M_1$. For the Jones projections e_K and e_{K_1} for the inclusions $K \subset M$ and $K_1 \subset M_1$, respectively, since $\mathcal{F}(e_K) = \lambda e_{K_1}$ for some scalar λ , we get the next theorem:

Theorem. If the depth of a given irreducible inclusion $N \subset M$ is 2, then an intermediate subfactor K of $N \subset M$ is normal in $N \subset M$ if and only if the depths of $N \subset K$ and $K \subset M$ are both 2.

The author [30] showed that if M is the crossed product $N \rtimes G$ of finite group G and $K = N \rtimes H$, then H is a normal subgroup of G if and only if $K \subset M \simeq K \subset K \rtimes F$ for some finite group F , i.e., the depth of $K \subset M$ is 2. Hence we see by the previous theorem that H is a normal subgroup of G if and only if K is a normal intermediate subfactor of $N \subset M$. Therefore our notion of normality for intermediate subfactors is an extension of that in the theory of finite groups.

2. PRELIMINARIES

2.1. Intermediate subfactors. We recall here some results for intermediate subfactors. Let $N \subset M$ be a pair of type II_1 factors. We denote by $\mathcal{L}(N \subset M)$ the set of all intermediate von Neumann subalgebras of $N \subset M$. The set $\mathcal{L}(N \subset M)$ forms a lattice under the two operations \vee and \wedge defined by

$$K_1 \vee K_2 = (K_1 \cup K_2)'' \text{ and } K_1 \wedge K_2 = K_1 \cap K_2.$$

If the relative commutant algebra $N' \cap M$ is trivial, then $\mathcal{L}(N \subset M)$ is exactly the lattice of intermediate subfactors for $N \subset M$. In fact for any $K \in \mathcal{L}(N \subset M)$, $\mathcal{Z}(K) = K' \cap K \subset N' \cap M = \mathbb{C}$. If M is the crossed product $N \rtimes_{\alpha} G$ for an outer action α of a finite group G , then it is well known that the intermediate subfactor lattice $\mathcal{L}(N \subset M)$ is isomorphic to the subgroup lattice $\mathcal{L}(G)$ (see [18], [19]). In [31] Y. Watatani proved the next theorem.

Theorem. Let $N \subset M$ be a pair of type II_1 factors. If $[M : N] < \infty$ and $N' \cap M = \mathbb{C}$, then $\mathcal{L}(N \subset M)$ is a finite lattice.

Later we were noted that this theorem was shown by S. Popa implicitly [23].

From now on we assume that $[M : N] < \infty$ and $N' \cap M = \mathbb{C}$. Let $N \subset M \subset M_1 \subset M_2$ be the Jones tower of $N \subset M$ obtained by iterating the basic extension. Let $e_N \in M_1$ and $e_M \in M_2$ be the Jones projections for $N \subset M$ and $M \subset M_1$, respectively. We denote by \mathcal{F} , Ocneanu's Fourier transform from $N' \cap M_1$ onto $M' \cap M_2$ i.e.,

$$\mathcal{F}(x) = [M : N]^{-\frac{3}{2}} E_{M'}^{N'}(x e_M e_N), \quad x \in N' \cap M_1,$$

where $E_{M'}^{N'}$ is the conditional expectation from N' onto M' . For $K \in \mathcal{L}(N \subset M)$, if e_K is the Jones projection for $K \subset M$, then e_K is an element of $N' \cap M_1$. In fact $K_1 = \langle M, e_K \rangle = J_M K' J_M \subset J_M N' J_M = M_1$ and hence $e_K \in K' \cap K_1 \subset N' \cap M_1$.

D. Bisch [1] and A. Ocneanu [21] gave the next characterization of intermediate subfactors in terms of Jones projections in $N' \cap M_1$.

Theorem. Let p be a projection in $N' \cap M_1$. There exists an intermediate subfactor $K \in \mathcal{L}(N \subset M)$ such that $p = e_K$ if and only if

$$(1) \ p \geq e_N,$$

$$(2) \ \mathcal{F}(p) = \lambda q \text{ for some } \lambda \in \mathbb{C} \text{ and some projection } q \in M' \cap M_2.$$

In this case, q is the Jones projection e_{K_1} for $K_1 \subset M_1$.

Lemma 2.1. *With the above notations, we have*

$$e_K = [K : N][M : N]E_{M_1}^{M_2}(e_M e_N e_{K_1}),$$

where $E_{M_1}^{M_2}$ is the trace preserving conditional expectation from M_2 onto M_1 .

Lemma 2.2. *Let K be an intermediate subfactor for $N \subset M$. Let $K \subset M \subset K_1 \subset K_2$ and $N \subset M \subset M_1 \subset M_2$ be the Jones towers for $K \subset M$ and $N \subset M$, respectively. If e_{K_1} is the Jones projection for $K_1 \subset M_1$, then there exists a $*$ -isomorphism φ of K_2 onto $e_{K_1}M_2e_{K_1}$ such that $\varphi(x) = xe_{K_1}$ for $x \in K_1$ and $\varphi(e_M^{K_1}) = e_M$, where $e_M^{K_1}$ and e_M are the Jones projections for $M \subset K_1$ and $M \subset M_1$, respectively.*

see for example [1], [27].

2.2. Finite dimensional Hopf C^* -algebras. In this subsection we recall some facts about finite dimensional Hopf C^* -algebras.

Let \mathbf{H} be a finite dimensional Hopf C^* -algebra with a comultiplication $\Delta_{\mathbf{H}}$ and an anti-pode $S_{\mathbf{H}}$. Let \mathbf{K} be a subHopf C^* -algebra of \mathbf{H} , i.e., \mathbf{K} is a $*$ -subalgebra of \mathbf{H} ,

$S_{\mathbf{H}}(\mathbf{K}) \subset \mathbf{K}$ and $\Delta_{\mathbf{H}}(\mathbf{K}) \subset \mathbf{K} \otimes \mathbf{K}$.

Lemma 2.3. *Define the subset \mathbf{K}^\perp of \mathbf{H}^* by*

$$\mathbf{K}^\perp = \{ f \in \mathbf{H}^* \mid (f, k) = 0, \forall k \in \mathbf{K} \},$$

where $(\cdot, \cdot) : \mathbf{H}^* \times \mathbf{H} \rightarrow \mathbb{C}$ is the dual pairing defined by $(f, h) = f(h)$, $f \in \mathbf{H}^*$, $h \in \mathbf{H}$.

Then \mathbf{K}^\perp is an ideal of \mathbf{H}^* .

Proof. Let g be an element of \mathbf{K}^\perp and f an element of \mathbf{H}^* . Then the element gf of \mathbf{H}^* is determined by the equation

$$(gf, h) = (g \otimes f, \Delta_{\mathbf{H}}(h)), \forall h \in \mathbf{H}.$$

By virtue of $\Delta_{\mathbf{H}}(\mathbf{K}) \subset \mathbf{K} \otimes \mathbf{K}$, we get

$$(gf, k) = (g \otimes f, \Delta_{\mathbf{H}}(k)) = 0, \forall k \in \mathbf{K}.$$

Therefore gf is an element of \mathbf{K}^\perp . Similarly, $fg \in \mathbf{K}^\perp$. \square

By the above lemma, there exists the central projection $p \in \mathbf{H}^*$ such that $\mathbf{K}^\perp = p\mathbf{H}^*$.

We put $e_{\mathbf{K}} = 1 - p$.

Proposition 2.4. *With the above notation, the reduced algebra $e_{\mathbf{K}}\mathbf{H}^*$ is the dual Hopf C^* -algebra of \mathbf{K} .*

Proof. Suppose that $k \in \mathbf{K}$ and $(y, k) = 0, \forall y \in e_{\mathbf{K}}\mathbf{H}^*$. Then

$$(f, k) = (e_{\mathbf{K}}f, k) + (pf, k) = (e_{\mathbf{K}}f, k) = 0, \forall f \in \mathbf{H}^*.$$

Therefore $k = 0$. Conversely, suppose that $y \in e_{\mathbf{K}}\mathbf{H}^*$ and $(y, k) = 0, \forall k \in \mathbf{K}$. Then $y \in \mathbf{K}^\perp \cap e_{\mathbf{K}}\mathbf{H}^* = \{0\}$. Hence the form $(\ , \)|_{e_{\mathbf{K}}\mathbf{H}^* \times \mathbf{K}}$ establishes a duality between \mathbf{K} and $e_{\mathbf{K}}\mathbf{H}^*$. So we can identify $e_{\mathbf{K}}\mathbf{H}^*$ with \mathbf{K}^* . Then for $y \in \mathbf{K}^*$ and $k_1, k_2 \in \mathbf{K}$, we have

$$(y, k_1 k_2) = (\Delta_{\mathbf{H}^*}(y), k_1 \otimes k_2) = (\Delta_{\mathbf{H}^*}(y)(e_{\mathbf{K}} \otimes e_{\mathbf{K}}), k_1 \otimes k_2).$$

Hence $\Delta_{\mathbf{K}^*}(y) = \Delta_{\mathbf{H}^*}(y)(e_{\mathbf{K}} \otimes e_{\mathbf{K}})$. Similarly, we have $S_{\mathbf{K}^*} = S_{\mathbf{H}^*}|_{\mathbf{K}^*}$ by the fact that

$$\overline{(y^*, k^*)} = (S_{\mathbf{H}^*}(y), k), \forall y \in \mathbf{K}^*, \forall k \in \mathbf{K}.$$

Therefore $e_{\mathbf{K}}\mathbf{H}^*$ is again a Hopf C^* -algebra with the dual algebra \mathbf{K} . \square

Theorem 2.5. *Let \mathbf{H} be a finite dimensional Hopf C^* -algebra. The number of subHopf C^* -algebras of \mathbf{H} is finite.*

Proof. By the above proposition, the map $\mathbf{K} \mapsto e_{\mathbf{K}}$ from the set of subHopf C^* -algebras of \mathbf{H} to central projections of \mathbf{H}^* is injective. Since the number of central projections of \mathbf{H}^* is finite, so is that of subHopf C^* algebras of \mathbf{H} . \square

Remark. Since every finite dimensional Hopf C^* -algebra (Kac algebra) admits an “outer” action on the AFD II_1 factor [34], the above theorem immediately follows from [31, Theorem 2.2].

Definition. Let \mathbf{H} be any Hopf algebra.

(1) The left adjoint action of \mathbf{H} on itself is given by

$$(ad_l h)(k) = \sum_{(h)} h_1 k (S_{\mathbf{H}}(h_2)), \text{ for all } h, k \in \mathbf{H}.$$

(2) The right adjoint action of \mathbf{H} on itself is given by

$$(ad_r h)(k) = \sum_{(h)} (S_{\mathbf{H}}(h_1)) k h_2, \text{ for all } h, k \in \mathbf{H}.$$

(3) A subHopf algebra \mathbf{K} of \mathbf{H} is called *normal* if both $(ad_l \mathbf{H})(\mathbf{K}) \subset \mathbf{K}$ and $(ad_r \mathbf{H})(\mathbf{K}) \subset \mathbf{K}$ hold. (See [17, pp. 33].)

The next proposition is useful later.

Proposition 2.6. *Let \mathbf{H} be a finite dimensional Hopf algebra with a counit $\varepsilon_{\mathbf{H}}$ and \mathbf{K} a subHopf algebra of \mathbf{H} . Then \mathbf{K} is normal if and only if $\mathbf{H}\mathbf{K}^+ = \mathbf{K}^+\mathbf{H}$, where $\mathbf{K}^+ = \mathbf{K} \cap \ker \varepsilon_{\mathbf{H}}$.*

See for a proof [17, pp. 35].

2.3. Bimodules. In this subsection we recall some facts about the bimodule calculus associated with an inclusion of type II_1 factors (see for example [21],[32]).

Let A, B, C be type II_1 factors and let $\alpha = {}_A H_B, \beta = {}_A K_B, \gamma = {}_B L_C$ be A - B , A - B and B - C Hilbert bimodules, respectively. We write $\alpha\gamma$ for the A - C Hilbert bimodule ${}_A H_B \otimes_B {}_B L_C$. We denote by $\langle \alpha, \beta \rangle$ the dimension of the space of A - B intertwiners from ${}_A H_B$ to ${}_A K_B$. The conjugate Hilbert space H^* of ${}_A H_B$ is naturally a B - A bimodule with B - A actions defined by

$$b \cdot \xi^* \cdot a = (a^* \xi b^*)^* \quad \text{for } a \in A \text{ and } b \in B,$$

where $\xi^* = \langle \cdot, \xi \rangle_H \in H^*$ for $\xi \in {}_A H_B$. We denote by $\bar{\alpha}$ the conjugate B - A Hilbert bimodule associated with α .

Proposition 2.7 (Frobenius reciprocity). *Let A, B, C be type II_1 factors, and $\alpha = {}_A H_B, \beta = {}_B K_C$ and $\gamma = {}_A L_C$ be Hilbert bimodules. Then*

$$\langle \alpha\beta, \gamma \rangle = \langle \alpha, \gamma\bar{\beta} \rangle = \langle \beta, \bar{\alpha}\gamma \rangle.$$

See for a proof [21], [32].

Example 2.8. Let M be a type II_1 factor with the normalized trace τ_M . As usual we let $L^2(M)$ be the Hilbert space obtained by completing M in the norm $\|x\|_2 = \sqrt{\tau_M(x^*x)}$, $x \in M$. Let $\eta : M \rightarrow L^2(M)$ be the canonical implementation. Let $J : L^2(M) \rightarrow L^2(M)$ be the modular conjugation defined by $J\eta(x) = \eta(x^*)$, $x \in M$. For $\theta \in \text{Aut}(M)$, we define ${}_M(L^2(M)_\theta)_M$, the M - M Hilbert bimodule, by

- (1) ${}_M(L^2(M)_\theta)_M = L^2(M)$ as a Hilbert space,
- (2) $x \cdot \xi \cdot y = xJ\theta(y)^*J\xi$, $x, y \in M$, $\xi \in L^2(M)$.

Then for $\theta, \theta_1, \theta_2 \in \text{Aut}(M)$ we have

$$\overline{{}_M(L^2(M)_\theta)_M} \simeq {}_M(L^2(M)_{\theta^{-1}})_M$$

$${}_M(L^2(M)_{\theta_1})_M \otimes_M {}_M(L^2(M)_{\theta_2})_M \simeq {}_M(L^2(M)_{\theta_1\theta_2})_M.$$

A bimodule $\alpha = {}_A H_B$ is called irreducible if $\langle \alpha, \alpha \rangle = 1$, i.e., $\text{End}_{A-B}({}_A H_B) \simeq \mathbb{C}$. If $\langle \alpha, \alpha \rangle < \infty$, $\alpha = {}_A H_B$, then we can get an A - B irreducible bimodule by cutting ${}_A H_B$ by a minimal projection in $\text{End}_{A-B}({}_A H_B)$.

Example 2.9. Let $N \subset M$ be an inclusion of type II_1 factors. We define the N - M bimodule ${}_N L^2(M)_M$ by actions $x \cdot \xi \cdot y = x J y^* J \xi$, $\xi \in L^2(M), x \in N, y \in M$. Then we can see that $\text{End}({}_N L^2(M)_M) \simeq N' \cap M$. In particular, if $N' \cap M = \mathbb{C}$, then ${}_N L^2(M)_M$ is an irreducible N - M bimodule.

The next lemma is well known.

Lemma 2.10. Let $N \subset M$ be a pair of type II_1 factors with finite index and M_1 the basic extension for the inclusion $N \subset M$. For $\theta \in \text{Aut}(N)$, ${}_N (L^2(M)_\theta)_N \simeq {}_N L^2(M)_N$ if and only if there exists a unitary $u \in M_1$ such that $u x u^* = \theta(x)$, for all $x \in N$, where ${}_N (L^2(M)_\theta)_N (= L^2(M) \text{ as a Hilbert space})$ is defined as in Example 2.8.

Example 2.11. Let $\gamma : G \rightarrow \text{Aut}(N)$ be an outer action of a finite group G on a type II_1 factor N . Let $M = N \rtimes_\gamma G$ be the crossed product and ρ the N - M bimodule ${}_N L^2(M)_M$ defined as in Example 2.9. If $\{\lambda_g \mid g \in G\}$ is a unitary implementation for the crossed product, then each element $x \in M$ is written in the form $x = \sum_{g \in G} x_g \lambda_g, x_g \in N$. This implies that the irreducible decomposition of $\rho \bar{\rho} = {}_N L^2(M)_N$ is

$$\bigoplus_{g \in G} {}_N (\overline{N \lambda_g}^{\|\cdot\|_2})_N \simeq \bigoplus_{g \in G} {}_N (L^2(N)_{\gamma_g})_N,$$

where ${}_N (L^2(N)_{\gamma_g})_N$ is the N - N bimodule as in Example 2.8.

3. DEFINITION OF NORMAL INTERMEDIATE SUBFACTORS

In this section, we shall introduce the notion of normality for intermediate subfactors and study its properties.

Let $N \subset M$ be a pair of type II_1 factors with $[M : N] < \infty$. Let $N \subset M \subset M_1 \subset M_2$ be the Jones tower of $N \subset M$, obtained by iterating the basic extensions. We denote by \mathcal{F} , Ocneanu's Fourier transform from $N' \cap M_1$ onto $M' \cap M_2$, i.e.,

$$\mathcal{F}(x) = [M : N]^{-\frac{3}{2}} E_{M'}^{N'}(x e_M e_N), \quad x \in N' \cap M_1,$$

where $E_{M'}^{N'}$ is the conditional expectation from N' onto M' .

Definition 3.1. Let K be an intermediate subfactor of $N \subset M$ and e_K the Jones projection for the inclusion $K \subset M$. Then we call that K is *normal* in $N \subset M$ if e_K and $\mathcal{F}(e_K)$ are elements of the centers of $N' \cap M_1$ and $M' \cap M_2$, respectively.

Lemma 3.2. *Let K be an intermediate subfactor for an irreducible inclusion $N \subset M$ of type II_1 factors with finite index. Let K_1 and M_1 be the basic extensions for $K \subset M$ and $N \subset M$, respectively. Then K is normal in $N \subset M$ if and only if K_1 is normal in $M \subset M_1$.*

Proof. Since $\mathcal{F}(e_K) = \lambda e_{K_1}$ for some $\lambda \in \mathbb{C}$, it is obvious by the definition. \square

Proposition 3.3. *Let $\alpha : G \rightarrow \text{Aut}(P)$ be an outer action of a finite group G on a type II_1 factor P and H a subgroup of G . Let M be the fixed point algebra $P^{(H, \alpha)}$ and N the fixed point algebra $P^{(G, \alpha)}$. For $K \in \mathcal{L}(N \subset M)$, there is a subgroup A of G*

such that $H \subset A \subset G$ and $K = P^{(A, \alpha)}$. Then K is a normal intermediate subfactor of $N \subset M$ if and only if $AgH = HgA$ for $\forall g \in G$. In particular, K is normal in $N \subset P$ if and only if A is normal subgroup of G .

Proof. Let $\{u_g \mid g \in G\}$ be unitary operators on $L^2(P)$ defined by $u_g \eta(x) = \eta(\alpha_g(x))$, $x \in P$, where $L^2(P)$ and η are defined as in Example 2.8. Let P_1 be the basic extension for $N \subset P$. Then $N' \cap P_1 = \{\sum_{g \in G} x_g u_g \mid x_g \in \mathbb{C}\} \simeq \mathbb{C}G$. Let e_M^P be the Jones projection for $M \subset P$. Then $e_M^P = \frac{1}{\#H} \sum_{h \in H} u_h$. Let M_1 be the basic extension for $N \subset M$. Then by Lemma 2.2,

$$N' \cap M_1 \simeq e_M^P(N' \cap P_1)e_M^P = \{\sum_{g \in G} \sum_{h, k \in H} x_g u_{h g k} \mid x_g \in \mathbb{C}\}.$$

Therefore

$$\begin{aligned} e_K^M \in \mathcal{Z}(N' \cap M_1) &\Leftrightarrow e_M^P e_K^P e_M^P (= e_K^P = \frac{1}{\#A} \sum_{a \in A} u_a) \in \mathcal{Z}(e_M^P(N' \cap P_1)e_M^P) \\ &\Leftrightarrow \sum_{a \in A} \sum_{h, k \in H} u_{a h g k} = \sum_{a \in A} \sum_{h, k \in H} u_{h g k a} \text{ for } \forall g \in G \\ &\Leftrightarrow AgH = HgA \text{ for } \forall g \in G. \end{aligned}$$

Since $M' \cap M_2$ is a commutative algebra, we get the the result. \square

We obviously have the dual version of this proposition by Lemma 3.2.

In [31] Y. Watatani introduced the notion of *quasi-normal intermediate subfactors* to study the modular identity for intermediate subfactor lattices.

Definition. Let $N \subset M$ be an inclusion of type II_1 factors with finite index and K an intermediate subfactor of $N \subset M$. Then K is *quasi-normal* (or *doubly commuting*)

if for any $L \in \mathcal{L}(N \subset M)$,

$$\begin{array}{ccc} K & \subset & K \vee L \\ \cup & & \cup \\ K \wedge L & \subset & L \end{array}$$

and

$$\begin{array}{ccc} K_1 & \subset & K_1 \vee L_1 \\ \cup & & \cup \\ K_1 \wedge L_1 & \subset & L_1 \end{array}$$

are commuting squares (see for example [5]), where K_1 and L_1 are the basic extension for $K \subset M$ and $L \subset M$, respectively.

Proposition 3.4. *Let $N \subset M$ be an irreducible inclusion of type II_1 factors with finite index. If K is a normal intermediate subfactor of $N \subset M$ then K is quasi-normal in $N \subset M$*

Proof. Suppose that the Jones projection e_K for $K \subset M$ is an element of the center of $N' \cap M_1$. Then for any intermediate subfactor L of $N \subset M$, since the Jones projection $e_K^{K \vee L}$ for $K \subset (K \vee L)$ is also a central projection in $K' \cap (K \vee L)_1$, we

have

$$\begin{array}{ccc} K & \subset & K \vee L \\ \cup & & \cup \\ K \wedge L & \subset & L \end{array}$$

is a commuting square. Similarly, if $\mathcal{F}(e_K)$ is an element of the center of $M' \cap M_2$, then

$$\begin{array}{ccc} K_1 & \subset & K_1 \vee L_1 \\ \cup & & \cup \\ K_1 \wedge L_1 & \subset & L_1 \end{array}$$

is a commuting square. Therefore if K is normal in $N \subset M$, then K is quasi-normal. \square

We have a characterization of normal intermediate subfactors in terms of bimodules. Let K be an intermediate subfactor of an irreducible inclusion $N \subset M$ of type II_1 factors with finite index. We note that e_K is in the center of $N' \cap M_1$ if and only if for any $T \in \text{End}({}_N L^2(M)_N)$, $TL^2(K) \subset L^2(K)$.

Proposition 3.5. *Let K be an intermediate subfactor for an irreducible inclusion $N \subset M$ of type II_1 factors with finite index. Let α be the N - K bimodule ${}_N L^2(K)_K$ and β the K - M bimodule ${}_K L^2(M)_M$. If ρ is the N - M bimodule $\alpha\beta = {}_N L^2(M)_M$, then K is normal in $N \subset M$ if and only if*

$$(1) \langle \alpha \bar{\alpha}, \rho \bar{\rho} \rangle = \langle \alpha \bar{\alpha}, \alpha \bar{\alpha} \rangle,$$

$$(2) \langle \bar{\beta} \beta, \bar{\rho} \rho \rangle = \langle \bar{\beta} \beta, \bar{\beta} \beta \rangle.$$

Proof. Since $\text{End}({}_N L^2(K)_N) = N' \cap \langle N, e_N^K \rangle \simeq e_K(N' \cap M_1)e_K$ by Lemma 2.2, if e_K is an element of the center of $N' \cap M_1$, then for any irreducible N - N bimodule σ contained in $\alpha \bar{\alpha}$, the multiplicity of σ in $\alpha \bar{\alpha}$ is equal to the multiplicity of σ in $\rho \bar{\rho}$. Therefore we have $\langle \alpha \bar{\alpha}, \rho \bar{\rho} \rangle = \langle \alpha \bar{\alpha}, \alpha \bar{\alpha} \rangle$. Conversely, suppose that e_K is not an element of the center of $N' \cap M_1$. Then there exist minimal projections $p \sim q$ in $(N' \cap M_1)$ such that $p \in e_K(N' \cap M_1)e_K$ and $q \notin e_K(N' \cap M_1)e_K$. Therefore we have $\langle \alpha \bar{\alpha}, \rho \bar{\rho} \rangle \neq \langle \alpha \bar{\alpha}, \alpha \bar{\alpha} \rangle$. And hence e_K is an element of the center of $(N' \cap M_1)$ if and only if $\langle \alpha \bar{\alpha}, \rho \bar{\rho} \rangle = \langle \alpha \bar{\alpha}, \alpha \bar{\alpha} \rangle$. Similarly, we can see that e_{K_1} is an element of the center of $(M' \cap M_2)$ if and only if $\langle \bar{\beta} \beta, \bar{\rho} \rho \rangle = \langle \bar{\beta} \beta, \bar{\beta} \beta \rangle$. Since $\mathcal{F}(e_K) = \lambda e_{K_1}$ for some $\lambda \in \mathbb{C}$, we get the result. \square

Theorem 3.6. *Let K be an intermediate subfactor for an irreducible inclusion $N \subset M$ of type II_1 factors with finite index. If the depths of $N \subset K$ and $K \subset M$ are both 2, then K is normal in $N \subset M$.*

Proof. Let α be the N - K bimodule ${}_N L^2(K)_K$ and β the K - M bimodule ${}_K L^2(M)_M$.

By the assumption, we have

$$\alpha \bar{\alpha} \alpha \simeq \underbrace{\alpha \oplus \alpha \oplus \cdots \oplus \alpha}_{[K:N] \text{ times}}$$

and

$$\overline{\beta\beta\beta} \simeq \underbrace{\overline{\beta} \oplus \overline{\beta} \oplus \cdots \oplus \overline{\beta}}_{[M:K] \text{ times}}.$$

And hence $\langle \alpha\overline{\alpha}, \alpha\overline{\alpha} \rangle = \langle \alpha\overline{\alpha}\alpha, \alpha \rangle = [K : N]$ and $\langle \overline{\beta\beta}, \overline{\beta\beta} \rangle = \langle \overline{\beta\beta\beta}, \overline{\beta} \rangle = [M : K]$ by Frobenius reciprocity. Since $N \subset M$ is irreducible, if ρ is the N - M bimodule ${}_N L^2(M)_M (= \alpha\beta)$, then $1 = \langle \rho, \rho \rangle = \langle \alpha\beta, \alpha\beta \rangle = \langle \overline{\alpha\alpha}, \overline{\beta\beta} \rangle$. And hence we have

$$\begin{aligned} \langle \alpha\overline{\alpha}, \rho\overline{\rho} \rangle &= \langle \alpha\overline{\alpha}, \alpha\beta\overline{\beta\alpha} \rangle = \langle \alpha\overline{\alpha}\alpha, \alpha\beta\overline{\beta} \rangle \\ &= [K : N] \langle \alpha, \alpha\beta\overline{\beta} \rangle = [K : N] \langle \overline{\alpha\alpha}, \overline{\beta\beta} \rangle = [K : N], \end{aligned}$$

i.e., $\langle \alpha\overline{\alpha}, \rho\overline{\rho} \rangle = \langle \alpha\overline{\alpha}, \alpha\overline{\alpha} \rangle$. Similarly, we have $\langle \overline{\beta\beta}, \overline{\rho\rho} \rangle = \langle \overline{\beta\beta}, \overline{\beta\beta} \rangle$. So we get the result by Proposition 3.5. \square

Proposition 3.7. *Let M_0, N_0, K be intermediate subfactors for an irreducible inclusion $N \subset M$ of type II_1 factors with finite index such that $N \subset N_0 \subset K \subset M_0 \subset M$. If K is normal in $N \subset M$, then K is also normal in $N_0 \subset M_0$.*

4. NORMAL INTERMEDIATE SUBFACTORS FOR DEPTH 2 INCLUSIONS

It is well-known that the crossed product of a finite dimensional Hopf C^* algebra (Kac algebra) is characterized by the depth 2 condition. In this section we study normal intermediate subfactors for depth 2 inclusions.

4.1. The action of $K' \cap K_1$ on M . Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index. Let $N \subset M \subset M_1 \subset M_2$ be the Jones tower for

$N \subset M$. We put $A = N' \cap M_1$ and $B = M' \cap M_2$. Then A and B are dual pair of Hopf C^* -algebras with a pairing

$$(a, b) = [M : N]^2 \tau(ae_M e_N b), \text{ for } a \in A \text{ and } b \in B,$$

where e_N and e_M are the Jones projections for $N \subset M$ and $M \subset M_1$, respectively.

Define a bilinear map $A \times M \rightarrow M$ (denoted by $a \odot x$) by setting

$$a \odot x = [M : N] E_M^{M_1}(axe_N),$$

for $x \in M$ and $a \in A$. This map is a left action of Hopf C^* algebra A and

$$N = M^A = \{ x \in M \mid a \odot x = \varepsilon(a)x, \forall a \in A \},$$

where $\varepsilon : A \rightarrow \mathbb{C}$ is the counit determined by $ae_N = \varepsilon(a)e_N$ (see [29]).

Proposition 4.1. *Let K be an intermediate subfactor of $N \subset M$ and K_1 the basic extension for $K \subset M$. We put $H = K' \cap K_1$. If a is an element of H , then*

$$[M : K] E_M^{K_1}(axe_K) = [M : N] E_M^{M_1}(axe_N), \forall x \in M.$$

This implies $K = M^H = \{ x \in M \mid a \odot x = \varepsilon(a)x, \forall a \in H \}$.

Proof. Since $e_K = \frac{[M:N]}{[M:K]} E_{K_1}^{M_1}(e_N)$ by [27], we have

$$\begin{aligned} [M : K] E_M^{K_1}(axe_K) &= [M : K] E_M^{K_1}(ax \frac{[M : N]}{[M : K]} E_{K_1}^{M_1}(e_N)) \\ &= [M : N] E_M^{K_1}(E_{K_1}^{M_1}(axe_N)) \\ &= [M : N] E_M^{M_1}(axe_N) \end{aligned}$$

for $\forall a \in H$ and $\forall x \in M$. \square

4.2. Hopf algebra structures on $K' \cap K_1$. Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index and K an intermediate subfactor of $N \subset M$. Then the depth of $K \subset M$ is not 2 in general. In this subsection we shall prove that if the depth of $K \subset M$ is 2, then $H = K' \cap K_1$ is a subHopf C^* algebra of $A = N' \cap M_1$.

By Lemma 2.2, there exists an isomorphism φ of K_2 onto $e_{K_1}M_2e_{K_1}$ such that $\varphi(x) = xe_{K_1}$ for $x \in K_1$ and $\varphi(e_M^{K_1}) = e_M$, where $K \subset M \subset K_1 \subset K_2$ is the Jones tower for the inclusion $K \subset M$ and $e_M^{K_1}$ is the Jones projection for $M \subset K_1$.

Lemma 4.2. *With the above notation, we have*

$$[M : K]^2 \tau(he_M^{K_1} e_K k) = [M : N]^2 \tau(he_M e_N \varphi(k))$$

for $\forall h \in H = K' \cap K_1$ and $\forall k \in M' \cap K_2$.

Lemma 4.3. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index and K an intermediate subfactor for $N \subset M$. Let $N \subset M \subset M_1 \subset M_2$ and $K \subset M \subset K_1 \subset K_2$ be the Jones towers for $N \subset M$ and $K \subset M$, respectively. If the depth of $K \subset M$ is 2, then for any $b \in M' \cap M_2$, there exist elements $\{x_i\}, \{y_i\}$ of $N' \cap M_1$ such that*

$$b = \sum_i x_i e_M y_i$$

and

$$\sum_i E_{K_1}^{M_1}(x_i) e_M E_{K_1}^{M_1}(y_i) \in (K' \cap K_1) e_M (K' \cap K_1),$$

where $E_{K_1}^{M_1}$ is the trace preserving conditional expectation from M_1 onto K_1 .

Proposition 4.4. *Suppose that $N \subset M$ is irreducible and the depth of $N \subset M$ is 2. Let K be an intermediate subfactor for $N \subset M$ and K_1 the basic extension for $K \subset M$. If the depth of $K \subset M$ is 2, then $H = K' \cap K_1$ is a subHopf algebra of $A = N' \cap M_1$.*

Theorem 4.5. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index and K an intermediate subfactor for $N \subset M$. Let $N \subset M \subset M_1 \subset M_2$ and $K \subset M \subset K_1 \subset K_2$ be the Jones towers for $N \subset M$ and $K \subset M$, respectively. Then the depth of $K \subset M$ is 2 if and only if e_{K_1} is an element of the center of $M' \cap M_2$, where e_{K_1} is the Jones projection for $K_1 \subset M_1$.*

Proof. Suppose that the depth of $K \subset M$ is 2. Then by the proof of Theorem 3.6, e_{K_1} is an element of the center of $M' \cap M_2$.

Conversely, suppose that e_{K_1} is an element of the center of $M' \cap M_2$. Then for any $h \in H = K' \cap K_1$, we have

$$\begin{aligned} (S_A(h), b) &= \overline{(h^*, b^*)} = [M : N]^2 \tau(b e_N e_M h) \\ &= [M : N]^2 \tau(e_{K_1} b e_{K_1} e_N e_M h) \\ &= (S_A(h), e_{K_1} b e_{K_1}) \quad \text{for } \forall b \in B = M' \cap M_2 \end{aligned}$$

and hence $S_A(H) \subset H$. Similarly, for any $h \in H$, we have

$$\begin{aligned} (\Delta_A(h), x \otimes y) &= (h, xy) = (h, e_{K_1} x e_{K_1} y e_{K_1}) \\ &= (\Delta_A(h), e_{K_1} x e_{K_1} \otimes e_{K_1} y e_{K_1}) \quad \text{for } \forall x, y \in M' \cap M_2, \end{aligned}$$

and hence $\Delta_A(H) \subset H \otimes H$. Therefore H is a subHopf algebra of $N' \cap M_1$. By Proposition 4.1, we have $K = M^H$. So the depth of $K \subset M$ is 2. \square

We obviously have the dual version by Lemma 3.2.

Remark. Later we noted by the referee that this theorem follows from the next characterization of depth 2 inclusions by bimodules: Let $N \subset M$ be an irreducible inclusion of type II_1 factors with finite index and ρ the N - M bimodule ${}_N L^2(M)_M$. Then the depth of $N \subset M$ is 2 if and only if for any irreducible bimodule ${}_N X_N \prec \rho \bar{\rho}$, $\dim_N X = \langle X, \rho \bar{\rho} \rangle$.

Theorem 4.6. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index and K an intermediate subfactor for $N \subset M$. Then K is a normal intermediate subfactor of $N \subset M$ if and only if the depths of $N \subset K$ and $K \subset M$ are both 2.*

Proof. This immediately follows from Theorem 4.5 and Lemma 3.2. \square

Theorem 4.7. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index and K an intermediate subfactor for $N \subset M$. Then K is a normal intermediate subfactor of $N \subset M$ if and only if $K' \cap K_1$ is a normal subHopf algebra of $N' \cap M_1$, where K_1 and M_1 are the basic extensions for $N \subset M$ and $K \subset M$, respectively.*

Proof. Suppose that K is a normal intermediate subfactor of $N \subset M$. Then $H =$

$K' \cap K_1$ is a subHopf algebra of $A = N' \cap M_1$ by Proposition 4.4. Let ε_H is a counit of H . Then $xe_K = \varepsilon_H(x)e_K$ for $x \in H$. Therefore $H^+ = H \cap \ker \varepsilon_H = H(1 - e_K)$. Since $(1 - e_K)$ is an element of the center of A by the assumption, we have $H^+A = AH^+$. Hence H is a normal subHopf algebra of A by Proposition 2.6. Conversely, we suppose that H is a normal subHopf algebra of A . Then by Proposition 4.4 and Theorem 4.5, e_{K_1} is element of the center of $M' \cap M_2$. Since $H^+ = H(1 - e_K)$, we have $(1 - e_K)A = A(1 - e_K)$ by Proposition 2.6. This implies e_K is an element of the center of $N' \cap M_1$ and hence K is a normal intermediate subfactor of $N \subset M$. \square

4.3. Lattices of normal intermediate subfactors. Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index. In this subsection we shall prove that the set of all normal intermediate subfactors of the inclusion $N \subset M$, denoted by $\mathcal{N}(N \subset M)$, is a sublattice of $\mathcal{L}(N \subset M)$. Moreover, $\mathcal{N}(N \subset M)$ is a modular lattice.

Lemma 4.8. *Let L and K be intermediate subfactors of $N \subset M$ and L_1 and K_1 the basic extensions for $L \subset M$ and $K \subset M$, respectively. Then the basic extension $(L \wedge K)_1$ for $(L \wedge K) \subset M$ is $L_1 \vee K_1$ and the basic extension $(L \vee K)_1$ for $(L \vee K) \subset M$ is $L_1 \wedge K_1$.*

We note that if we denote by e_A the Jones projection for $A \subset M$, then for $L, K \in \mathcal{L}(N \subset M)$, we have $e_{L \wedge K} = e_L \wedge e_K$. But $e_{L \vee K} \neq e_L \vee e_K$ in general (see [27]).

Theorem 4.9. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index. Then the set of all normal intermediate subfactors $\mathcal{N}(N \subset M)$ is a sublattice of $\mathcal{L}(N \subset M)$*

Proof. Let L and K be normal intermediate subfactors of $N \subset M$. Since e_L and e_K are elements of the center of $N' \cap M_1$ by the assumption, we have $e_{L \wedge K} = e_L \wedge e_K \in \mathcal{Z}(N' \cap M_1)$ by the above argument. Observe that

$$(L \vee K)' \cap (L \vee K)_1 = (L' \cap L_1) \cap (K' \cap K_1).$$

Since $L' \cap L_1$ and $K' \cap K_1$ are invariants under the left and right adjoint action of $N' \cap M_1$ (see Definition in 2.2), so is $(L \vee K)' \cap (L \vee K)_1$. Therefore we can see that $(L \vee K)' \cap (L \vee K)_1$ is a normal subHopf algebra $N' \cap M_1$ by the definition. Since $L \vee K$ is a normal intermediate subfactor of $N \subset M$ by Theorem 4.7, we have $e_{L \vee K} \in \mathcal{Z}(N' \cap M_1)$. Applying the same argument for the dual inclusion $M \subset M_1$, we conclude that $L \wedge K$ and $L \vee K$ are normal intermediate subfactors of $N \subset M$. \square

Corollary 4.10. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors. Then $\mathcal{N}(N \subset M)$ is a modular lattice.*

Proof. This immediately follows from Proposition 3.4, Theorem 4.9 and [31, Theorem 3.9]. \square

Theorem 4.11. *Let $N \subset M$ be an irreducible, depth 2 inclusion of type II_1 factors with finite index. Then every maximal chain from M to N in $\mathcal{N}(N \subset M)$ has the*

same length, i.e., for $A_i (i = 1, 2, \dots, m)$, $B_j (j = 1, 2, \dots, n) \in \mathcal{N}(N \subset M)$, if

$$M = A_0 > A_1 > \dots > A_m = N$$

and

$$M = B_0 > B_1 > \dots > B_n = N,$$

then $m = n$, where $X > Y$ means $X \supset Y$ and $X \supseteq K \supseteq Y$, implies $K = X$ or $K = Y$ for $X, Y, K \in \mathcal{N}(N \subset M)$.

Proof. Since we have the Jordan-Dedekind chain condition holding in modular lattices, this immediately follows from the previous corollary. (see for example [6].) \square

Example 4.12. We denote by S_n the symmetric group on n letters, x_1, x_2, \dots, x_n and $\sigma = (1, 2, 3, \dots, n)$ the element of S_n with order n and $\langle \sigma \rangle$ the cyclic group generated by σ . Let $\gamma : S_n \rightarrow \text{Aut}(P)$ be an outer action of S_n on a type II_1 factor P and let $N = P^{\gamma_\sigma} \subset M = P \rtimes_\gamma S_{n-1}$. Then we can see that $S_n = S_{n-1} \langle \sigma \rangle = \langle \sigma \rangle S_{n-1}$ and $S_{n-1} \cap \langle \sigma \rangle = \{e\}$. Therefore the depth of $N \subset M$ is 2 (see [26, 33]). We put $K = P \rtimes_\gamma A_{n-1}$, where A_{n-1} is the alternating group consists of the even permutations on x_1, x_2, \dots, x_{n-1} . If n is odd, then the length of $\mathcal{N}(N \subset M)$ is 3 and if n is even, then that is 2.

REFERENCES

1. D. Bisch, *A note on intermediate subfactors*, Pacific J. Math. **163** (1994), 201–216.
2. D. Bisch and U. Haagerup, *Composition of subfactors: new examples of infinite depth subfactors*, preprint.

3. M. Choda and H. Kosaki, *Strongly outer actions for an inclusion of factors*, J. Funct. Anal. **122** (1994), 315–332.
4. D. Evans and Y. Kawahigashi, *Orbifold subfactors from Hecke algebras*, Comm. Math. Phys. **165** (1994), 445–484.
5. F. Goodman, P. de la Harpe, and V. F. R. Jones, *Coxeter graph and tower of algebras*, vol. 14, MSRI Publ, Springer-Verlag, 1989.
6. M. Hall, Jr., *The theory of groups*, The Macmillan Company, New York, 1959.
7. M. Izumi, *Application of fusion rules to classification of subfactors*, Publ. RIMS, Kyoto Univ. **27** (1991), 953–994.
8. ———, *Goldman's type theorem for index 3*, Publ. RIMS, Kyoto Univ. **28** (1992), 833–843.
9. M. Izumi and Y. Kawahigashi, *Classification of subfactors with the principal graph $D_n^{(1)}$* , J. Funct. Anal. **112** (1993), 257–286.
10. V. F. R. Jones, *Index for subfactors*, Invent.math. **72** (1983), 1–25.
11. Y. Kawahigashi, *On flatness of Ocneanu's connection on the dynkin diagrams and classification of subfactors*, J. Funct. Anal. **127** (1995), 63–107.
12. H. Kosaki, *Characterization of crossed product (properly infinite case)*, Pacific J. Math. **137** (1989), 159–167.
13. H. Kosaki and R. Longo, *A remark on the minimal index of subfactors*, J. Funct. Anal. **107** (1992), 458–470.
14. P. H. Loi, *On the theory of index and type III factors*, Ph.D. thesis, Pennsylvania State Univ., 1988.
15. R. Longo, *Index of subfactors and statistics of quantum fields I*, Commun. Math. Phys. **126** (1989), 217–247.
16. ———, *Index of subfactors and statistics of quantum fields II*, Commun. Math. Phys. **130**

- (1990), 285–390.
17. S. Montgomery, *Hopf algebras and their actions on rings*, CBMS series number 82, 1992.
 18. M. Nakamura and Z. Takeda, *On the fundamental theorem of the Galois theory for finite factors*, Japan Acad. **36** (1960), 313–318.
 19. ———, *A Galois theory for finite factors*, Proc. Japan Acad. **36** (1960), 258–260.
 20. A. Ocneanu, *Quantized groups, string algebras and galois theory for algebras*, Operator Algebras and Applications, vol.2, London Math. Soc. Lecture Note Series Vol.136, Cambridge Univ. Press, 1988, pp. 119–172.
 21. ———, *Quantum symmetry, differential geometry of finite graphs, and classification of subfactors*, 1991, Univ. of Tokyo Seminary Notes.
 22. P. Pimsner and S. Popa, *Entropy and index for subfactors*, Ann. Sci. Ecole Norm. Sup. **19** (1986), 57–106.
 23. S. Popa, *Correspondences*, preprint.
 24. ———, *Classification of subfactors: the reduction to commuting squares*, Invent. Math. **101** (1990), 19–43.
 25. ———, *Classification of amenable subfactors of type II*, Acta Math. **172** (1994), 163–255.
 26. T. Sano, *Commuting co-commuting squares and finite dimensional Kac algebras*, to appear in Pacific J. Math.
 27. T. Sano and Y. Watatani, *Angles between two subfactors*, J. Operator Theory **32** (1994), 209–242.
 28. Y. Sekine, *Connections associated with finite-dimensional Kac algebra actions*, Kyushu J. Math. **49** (1995), 253–269.
 29. W. Szymański, *Finite index subfactors and Hopf algebra crossed products*, Proc. Amer. Math. Soc. **120** (1994), 519–528.

- 30. T. Teruya, *A characterization of normal extensions for subfactors*, Proc. Amer. Math. Soc. **120** (1994), 781–783.
- 31. Y. Watatani, *Lattices of intermediate subfactors*, J. Funct. Anal., to appear.
- 32. S. Yamagami, *A note on Ocneanu's approach to Jones' index theory*, Internat. J. of Math. **4** (1993), 859–871.
- 33. ———, *Vector bundles and bimodules*, Quantum and Non-Commutative Analysis, Kluwer Academic, 1993, pp. 321–329.
- 34. T. Yamanouchi, *Construction of an outer action of a finite-dimensional Kac algebra on the AFD factor of type II_1* , Inter. J. Math. **4** (1993), 1007–1045.